

Magneto-acoustic rotation of transverse waves in $^3\text{He-B}$

J. A. Sauls^{a,1}, Y. Lee^b, T.M. Haard^a, W.P. Halperin^a

^a*Department of Physics and Astronomy, Northwestern University, Evanston IL 60208 U.S.A.*

^b*present address: Department of Physics, Stanford University, Stanford CA 94305 U.S.A.*

Abstract

In superfluid $^3\text{He-B}$, the off-resonant coupling of the $J = 2^-$, $M = \pm 1$ order parameter collective modes to transverse current excitations stabilizes propagating transverse waves with low damping for frequencies above that of the $J = 2^-$ modes. Right- (RCP) and left circularly polarized (LCP) transverse modes are degenerate in zero field; however, a magnetic field with $\mathbf{H} \parallel \mathbf{q}$ lifts this degeneracy giving rise to the acoustic analog of circular birefringence and an acoustic Faraday effect for linearly polarized transverse sound waves [1]. We present theoretical results for the temperature, pressure and field dependence of the Faraday rotation angle, and compare the theory with recent measurements [2]. The analysis provides a direct measurement of the Landé g-factor for the $J = 2^-$ modes, and new information on the magnitude of f-wave pairing correlations in $^3\text{He-B}$.

Keywords: Superfluid ^3He ; Transport; Acoustics; Collective Modes; Nuclear Zeeman Effect

The dispersion relation for RCP(+) and LCP(-) transverse current modes in $^3\text{He-B}$ is given by [1]

$$\left(\frac{\omega}{q_{\pm} v_f}\right)^2 = \Lambda_n + \Lambda_s \frac{\omega^2}{(\omega + i\Gamma)^2 - \Omega_{\pm}^2(T, \omega, H)}, \quad (1)$$

with $\Lambda_n = \frac{F_1^s}{15} \rho_n(\omega, T)$ and $\Lambda_s = \frac{2F_1^s}{75} \rho_s(\omega, T)$. The restoring forces are provided by quasiparticle excitations ($\sim \rho_n$), and pair excitations ($\sim \rho_s$), with $\rho_s + \rho_n = 1$. The condensate term dominates at low temperature ($\rho_s \simeq 1$), and is anomalously large when the sound frequency is near a resonant frequency of the $J = 2^-$, $M = \pm 1$ collective modes, *i.e.* when

$$D_{\pm}^2(\omega, H, T) = (\omega + i\Gamma)^2 - \Omega_{\pm}^2(\omega, H, T) \gtrsim 0. \quad (2)$$

The frequencies of the $M = \pm 1$ modes include the Zeeman splitting,

$$\Omega_{\pm}^2 = \omega_{isq}^2(T) \pm 2g(T)\omega\omega_L(H, T), \quad (3)$$

where $\omega_{isq}(T)$ is the frequency of the $J = 2^-$ modes in zero field, $g(T)$ is the g-factor for the $J = 2^-$ modes and $\omega_L(H, T) = \gamma_{\text{eff}} H$ is the effective Larmor frequency that determines the linear Zeeman splitting of the $J = 2$ multiplets [3].

Consider a linearly polarized transverse current excitation with frequency ω propagating in the z -direction. The RCP and LCP modes propagate in the bulk with different phase velocities,

$$\mathbf{J}(\omega, z) = \frac{J}{\sqrt{2}} e^{iq_+(\omega)z} \hat{\mathbf{e}}_+ + \frac{J}{\sqrt{2}} e^{iq_-(\omega)z} \hat{\mathbf{e}}_-, \quad (4)$$

where $\hat{\mathbf{e}}_{\pm} = (\hat{\mathbf{x}} \pm i\hat{\mathbf{y}})/\sqrt{2}$ are the polarization vectors for RCP and LCP current modes. The re-

¹ Corresponding author. E-mail: sauls@nwu.edu

sponse corresponds to a pure Faraday rotation of the polarization if the phase velocities are real, which is the case at very low temperatures ($T \ll T_c$) and frequencies above the collective mode resonances ($|\omega - \Omega_{\pm}(T)| \gg \Gamma$). The transverse wave propagates with the average phase velocity, $\bar{c} = 2\omega/(q_+ + q_-)$, while the polarization *rotates* with a spatial period, $\Lambda_H(\omega, T) = 4\pi/|q_+ - q_-|$.

Near the $M = +1$ mode crossing ($\omega = \Omega_+(T_+)$) the temperature dependence of the Faraday rotation period is dominated by $q_+(T) \simeq q_f \sqrt{s(T/T_+ - 1)}$, where $q_f = \omega/v_f$ and $s \equiv 2|\Omega_+|T_+/\Lambda_s\Omega_+$ are evaluated at $T = T_+$. The field dependence of Λ_H originates from the Zeeman splitting of the $M = \pm 1$ modes, and appears through the shift in the wavenumber of the LCP mode. Using $D_-^2 = D_+^2 + 4g(T)\omega\omega_L(T, H)$ we can write, $q_- \simeq q_f \sqrt{s(T/T_+ - 1) + B/B_+}$, where we set $g = g(T_+)$ for low fields, $g\omega_L \ll \Omega_+$, and $T \gtrsim T_+$. The formula for $q_-(T)$ also simplifies because $\Lambda_n D_-^2(T) \ll \Lambda_s \omega^2$. The field dependence is determined by the field scale, $B_+ \equiv \Lambda_s \Omega_+/4g\gamma_{\text{eff}}$, where γ_{eff} is the effective gyromagnetic ratio. Thus, near the mode crossing the Faraday rotation period simplifies to

$$\Lambda_H = \frac{\lambda_f}{\sqrt{\alpha + \beta} - \sqrt{\alpha}}, \quad (5)$$

where $\lambda_f = 4\pi v_f/\Omega_+$, $\alpha = s(T/T_+ - 1)$, and $\beta = H/B_+$. Except for temperatures very close to the collective mode resonance or for very low fields; i.e. for $s(T/T_+ - 1) > H/B_+$, Λ_H scales inversely with H and as the square root of the reduced temperature, $\Lambda_H \simeq K \sqrt{T/T_+ - 1}/H$, where $K \simeq \xi\Omega(\pi v_f/\gamma_{\text{eff}}g) \sqrt{\frac{2}{25}(m^*/m - 1)\rho_s/\rho}$ is determined by the effective mass, superfluid density, gyromagnetic ratio, Landé g-factor for the modes, and the slope of the collective mode frequency, which is approximately $\xi\Omega = 8|\Omega'_+|T_+/\Omega_+ \simeq 1$ at $T \simeq 0.44T_c$. Recent experimental measurements of the Faraday rotation angle [2] confirm this scaling behavior. A quantitative comparison between the theory and experimental measurements of the Faraday rotation period as a function of temperature, pressure and field can be made on the basis of Eq. 5.




Fig. 1. Comparison between the theoretical and experimental results for the temperature dependence of the Faraday rotation period for a field of $H = 100$ Gauss, frequency of $\omega/2\pi = 82.26$ MHz at pressures, $p = 4.31$ bar (red), $p = 4.42$ bar (blue), and $p = 4.52$ bar (green) [2].

Fig. 1 shows the comparison between the theoretical and experimental results for the temperature dependence of the Faraday rotation period scaled to $H = 100$ Gauss for three pressures near 4 bar. The inputs to the theoretical calculation are: the Fermi liquid parameters, T_c , v_f , F_1^s , F_0^s [3], the mode data, T_+/T_c and ω , and the gyromagnetic ratio for ^3He . The calculated parameters that enter Eq. 5 are the effective Larmor frequency, $\omega_L = \gamma_{\text{eff}}H$, the condensate response, $\rho_s(\omega, T_+)$ and the slope of the mode frequency, $|\Omega'_+|T_+$. The remaining open parameter is the Landé g-factor which determines the vertical scale for Λ_H ; the fit to the data gives $g = 0.02 \pm 0.002$, which is in agreement with theoretical calculations of the g-factor that include attractive f-wave pairing correlations in $^3\text{He-B}$ [4].

References

- [1] G.F. Moores, J.A. Sauls, JLTTP 91 (1993) 13.
- [2] Y. Lee et al., Nature 400 (1999) 431.
- [3] W. P. Halperin, E. Varoquaux, in *Helium Three* (Halperin, W. P. and Pitaevskii, L. P., eds.), 353, Amsterdam: Elsevier Science Publishers, 1990.
- [4] J. A. Sauls, J. W. Serene, Phys. Rev. Lett. 49 (1982) 1183.